Neuroscience Letters, 182 (1993) 307–310
Elsevier Ireland Ltd.

AUTHOR INDEX

Abe, M., see Tsuzuki, K., 151
Agid, Y., see Villares, J., 59
Ågren, H., see Mathé, A.A., 138
Alleva, E., see Betancur, C., 147
Anderton, B.H., see Willis, C.L., 159
Anglade, F., Bizot, J.-C., Dodd, R.H., Baudoin, C. and Chapouthier, G., Opposite effects of cholinergic agents and benzodiazepine receptor ligands in a passive avoidance task in rats, 247
Angus-Leppan, H., Olausson, B., Boers, P. and Lambert, G.A., Convergence of afferents from superior sagittal sinus and tooth pulp on cells in the upper cervical spinal cord of the cat, 275
Arai, N., Papp, M.I. and Lantos, P.L., New observation on ubiquitinated neurons in the cerebral cortex of multiple system atrophy (MSA), 197
Araki, K., see Hirano, T., 172
Aubineau, P., see Mathiau, P., 133
Avoli, M., see Barbarosie, M.T., 208
Azev, O.A., see Koulchitsky, S.V., 129

Baba, A., Yamamoto, T., Yamamoto, H., Suzuki, T. and Moroji, T., Effects of the major metabolite of phencyclidine, the *trans* isomer of 4-phenyl-4-(1-piperidinyl)cyclohexanol, on [^{3}H]N-(1-[2-thienyl]cyclohexyl)-3,4-piperidine ([^{3}H]TCP) binding and [^{3}H]dopamine uptake in the rat brain, 119
Badia, A., see Clos, M.V., 125
Bakalara, N., see Mathiau, P., 133
Ban, Y., see Okamura, H., 181
Barbarosie, M.T., Nagao, T. and Avoli, M., Control of 4-aminopyridine-induced synchronous activity by adenosine A1 and μ-opioid receptor agonists in adult rat hippocampus, 208
Barrionuevo, G., see Chinestra, P., 177
Baubet, V., see Sermet, E., 213
Baudoin, C., see Anglade, F., 247
Ben-Ari, Y., see Chinestra, P., 177
Bentivoglio, M., see Peng, Z.-C., 104
Betancur, C., Dell'Omo, G. and Alleva, E., Magnetic field effects on stress-induced analgesia in mice: modulation by light, 147
Bielavska, E. and Krivanek, J., Intracerebral injection of polymyxin B blocks the acquisition of conditioned taste aversion in rats, 239
Bijak, M., see Rohrbacher, J., 95
Bizot, J.-C., see Anglade, F., 247
Blanco, R., see Ferrer, I., 77
Blandl, T., see Kardos, J., 73
Bobillier, P., see Sermet, E., 213
Boers, P., see Angus-Leppan, H., 275
Bowenkamp, K., see Hoffer, B.J., 107
Brickley, S.G., Keating, M.J. and Grant, S., Experience-dependent mechanism of binocular map plasticity in *Xenopus*: incongruent connections are masked by retinal input, 13
Brown, G.C., see Cady, E.B., 201
Bunker, M.C., see Hall, J.C., 222
Buselmaier, W., see Seipp, S., 91
Butterworth, R.F., see Rao, V.L.R., 69

Cady, E.B., Lorek, A., Penrice, J., Wylezinska, M., Cooper, C.E., Brown, G.C., Owen-Reece, H., Kirkbride, V., Wyatt, J.S., Osmund, E. and Reynolds, R., Brain-metabolite transverse relaxation times in magnetic resonance spectroscopy increase as adenosine triphosphate depletes during secondary energy failure following acute hypoxia–ischaemia in the newborn piglet, 201
Candy, J.M., see Ince, P.G., 87
Carcereri De Prati, A., see Menegazzi, M., 295
Cash, D.J., see Kardos, J., 73
Castanon, N., see Chaouloff, F., 217
Chaouloff, F., Castanon, N. and Mormède, P., Paradoxical differences in animal models of anxiety among the Roman rat lines, 217
Chapouthier, G., see Anglade, F., 247
Chen, X., see Hansen, L.A., 63
Chinestra, P., Diabira, D., Urban, N.N., Barrionuevo, G. and Ben-Ari, Y., Major differences between long-term potentiation and ACPD-induced slow onset potentiation in hippocampus, 177
Ciofi, P., Krause, J.E., Prins, G.S. and Mazzuca, M., Presence of nuclear androgen receptor-like immunoreactivity in neurokinin B-containing neurons of the hypothalamic arcuate nucleus of the adult male rat, 193
Clark, R.B., Kinsberg, E.R. and Giles, W.R., Histochemical localization of nitric oxide synthase in the Bullfrog intracardiac ganglion, 255
Clifton, G., see Yang, K., 287
Clifton, G., see Yang, K., 291
Clos, M.V., Garcia Sanz, A., Sabriá, J., Pastor, C. and Badia, A., Differential contribution of L- and N-type calcium channels on rat hippocampal acetylcholine release, 125
Cooper, C.E., see Cady, E.B., 201
Crook, R., see Hardy, J., 1
Ćulić, M., Šaponjić, J., Janković, B. and Rakić, L., Amphetamine and haloperidol modulatory effects on Purkinje cell activity and on EEG power spectra in the acute rat model of epilepsy, 259

Dalery, J., see Sermet, E., 213
Davies, A.M. and Rosenthal, A., Neurons from mouse embryos with a null mutation in the tumour suppressor gene p53 undergo normal cell death in the absence of neurotrophins, 112
Dell'Omo, G., see Betancur, C., 147
Démonet, J.-F., Price, C., Wise, R. and Frackowiak, R.S.J., Differential activation of right and left posterior sylvian regions by semantic and phonological tasks: a positron-emission tomography study in normal human subjects, 25
Diabira, D., see Chinestra, P., 177
Dodd, R.H., see Anglade, F., 247

Ehmann, W.D., see Ince, P.G., 87

Faucheux, B., see Villares, J., 59
Faucon Biguet, N., see Lavergne, A., 167
Faustinella, F., see Yang, K., 291
Faustinella, F., see Yang, K., 287
Ferrer, I., see Soriano, M.A., 205
Ferrer, I., Tortosa, A., Blanco, R., Martín, F., Serrano, T., Planas, A. and Macaya, A., Naturally occurring cell death in the developing cerebral cortex of the rat. Evidence of apoptosis-associated internucleosomal DNA fragmentation, 77
Feyerabend, B., Malz, C.R. and Meyer, D.L., Birds that feed-on-the-wing have few isthmo-optic neurons, 66
Fink-Jensen, A. and Kristensen, P., Effects of typical and atypical neuroleptics on Fos protein expression in the rat forebrain, 115

Fortin, M. and Parent, A., Patches in the striatum of squirrel monkeys are enriched with calretinin fibers but devoid of calretinin cell bodies, 51
Frackowiak, R.S.J., see Démonet, J.-F., 25
Frain, O., see Lavergne, A., 167
Fujii, N., see Tsuzuki, K., 151
Fujiwara, F., Hirai, H., Tamaki, N. and Okada, Y., Adenosine enhances neuronal damage during deprivation of oxygen and glucose in guinea pig superior collicular slices, 283
Fukatsu, R., see Tsuzuki, K., 151
Fukui, Y., see Hisano, S., 80
Furukawa, T., see Ogawa, S., 123

Gadient, R.A. and Otten, U., Identification of interleukin-6 (IL-6)-expressing neurons in the cerebellum and hippocampus of normal adult rats, 243
Galasko, D., see Hansen, L.A., 63
Garcia Sanz, A., see Clos, M.V., 125
Garner, C.C., see Huang, D.Y., 55
Gass, P., Schröder, H., Prior, P. and Kiessling, M., Constitutive expression of heat shock protein 90 (HSP90) in neurons of the rat brain, 188
Gay, N., see Sermet, E., 213
Gerhardt, G.A., see Hoffer, B.J., 107
Gerig, G., see Kikinis, R., 7
Giaume, M., see Sermet, E., 213
Giles, W.R., see Clark, R.B., 255
Goedert, M., see Huang, D.Y., 55
Gourine, A.V., see Koulchitsky, S.V., 129
Grant, S., see Brickley, S.G., 13
Grassi Zucconi, G., see Menegazzi, M., 295
Grönroos, M., Naukkarinen, H. and Pertovaara, A., Capsaicin-induced central facilitation of a sympathetic vasoconstrictor response to painful stimulation in humans, 163
Guerrero, J.M., see Harmouch, A., 303
Guibert, B., see Lavergne, A., 167

Haimson, J., see Kikinis, R., 7
Hall, J.C. and Bunker, M.C., Acetylcholinesterase staining in the auditory brainstem nuclei of the leopard frog, *Rana pipiens*, 222
Hansen, L.A., Galasko, D., Samuel, W., Xia, Y., Chen, X. and Saitoh, T., Apolipoprotein-E ε-4 is associated with increased neurofibrillary pathology in the Lewy body variant of Alzheimer's disease, 63
Hao, J.-X., see Xu, X.-J., 263
Hao, J.-X., Xu, X.-J. and Wiesenfeld-Hallin, Z., Intrathecal γ-aminobutyric acid$_B$ ($GABA_B$) receptor antagonist CGP 35348 induces hypersensitivity to mechanical stimuli in the rat, 299
Hardy, J., Crook, R., Prihar, G., Roberts, G., Raghavan, R. and Perry, R., Senile dementia of the Lewy body type has an apolipoprotein E ε4 allele frequency intermediate between controls and Alzheimer's disease, 1
Harmouch, A., Guerrero, J.M. and Osuna, C., Different sensitivity of rat pineal *N*-acetyltransferase to α- and β-adrenergic receptor agonists during development: in vitro studies, 303
Hayes, R.L., see Yang, K., 287
Hayes, R.L., see Yang, K., 291
Heiss, W.-D., see Herholz, K., 47
Herholz, K., Pietrzyk, U., Karbe, H., Würker, M., Wienhard, K. and Heiss, W.-D., Individual metabolic anatomy of repeating words demonstrated by MRI-guided positron emission tomography, 47
Hirai, H., Ikeuchi, Y. and Okada, Y., The contribution of PKA to the excitatory mechanism of adenosine in guinea pig superior colliculus slices, 33
Hirai, H., see Fujiwara, F., 283
Hiramatsu, K. and Ohshima, K., Colocalization of NADPH-diaphorase with neuropeptides in the intrapancreatic neurons of the chicken, 37
Hirano, T., Kasono, K., Araki, K., Shinozuka, K. and Mishina, M., Involvement of the glutamate receptor δ2 subunit in the long-term depression of glutamate responsiveness in cultured rat Purkinje cells, 172
Hirsch, E.C., see Villares, J., 59
Hisano, S., Uemura, N., Fukui, Y., Miki, M. and Zhang, R., Induction of Fos-like immunoreactivity in the rat hypothalamus by an endogenous feeding suppressant (2-buten-4-olide), 80
Hoffer, B.J., Hoffman, A., Bowenkamp, K., Huettl, P., Hudson, J., Martin, D., Lin, L.-F.H. and Gerhardt, G.A., Glial cell line-derived neurotrophic factor reverses toxin-induced injury to midbrain dopaminergic neurons in vivo, 107
Hoffman, A., see Hoffer, B.J., 107
Hokama, H., see Kikinis, R., 7
Hrabovszky, E. and Liposits, Z., Adrenergic innervation of dopamine neurons in the hypothalamic arcuate nucleus of the rat, 143
Hsu, K.-S., Modulation of the nicotinic acetylcholine receptor channels by spermine in *Xenopus* muscle cell culture, 99
Huang, D.Y., Goedert, M., Jakes, R., Weisgraber, K.H., Garner, C.C., Saunders, A.M., Pericak-Vance, M.A., Schmechel, D.E., Roses, A.D. and Strittmatter, W.J., Isoform-specific interactions of apolipoprotein E with the microtubule-associated protein MAP2c: implications for Alzheimer's disease, 55
Hudson, J., see Hoffer, B.J., 107
Huettl, P., see Hoffer, B.J., 107
Huganir, R.L., see Mei, L., 21

Ibata, Y., see Okamura, H., 181
Ikarashi, Y., see Takahashi, A., 235
Ikeuchi, Y., see Hirai, H., 33
Imamura, K., see Takechi, H., 279
Ince, P.G., Shaw, P.J., Candy, J.M., Mantle, D., Tandon, L., Ehmann, W.D. and Markesbery, W.R., Iron, selenium and glutathione peroxidase activity are elevated in sporadic motor neuron disease, 87
Inouye, S.-I.T., see Okamura, H., 181
Ishimaru, H., see Takahashi, A., 235
Itzhak, Y. and Norenberg, M.D., Attenuation of ammonia toxicity in mice by PK 11195 and pregnenolone sulfate, 251
Iwakiri, H., Oka, T. and Mori, S., Multi-segmental and generalized suppression of postural muscle tone evoked by stimulating the dorsal tegmental field and the medial pontine reticular formation in acute decerebrate cats, 83

Jakes, R., see Huang, D.Y., 55
Janković, B., see Ćulić, M., 259
Javoy-Agid, F., see Villares, J., 59
Jolesz, F.A., see Kikinis, R., 7

Kakiuchi, T., see Takechi, H., 279
Kamegai, M., see Konishi, Y., 271
Kampfl, A., see Yang, K., 291
Kampfl, A., see Yang, K., 287
Kanemasa, K., see Okamura, H., 181
Karbe, H., see Herholz, K., 47
Kardos, J., Kovács, I., Blandl, T. and Cash, D.J., Modulation of GABA flux across rat brain membranes resolved by a rapid quenched incubation technique, 73
Kasono, K., see Hirano, T., 172

Katsuragi, T., see Ogawa, S., 123
Keating, M.J., see Brickley, S.G., 13
Kiessling, M., see Gass, P., 188
Kikinis, R., Shenton, M.E., Gerig, G., Hokama, H., Haimson, J., O'Donnell, B.F., Wible, C.G., McCarley, R.W. and Jolesz, F.A., Temporal lobe sulco-gyral pattern anomalies in schizophrenia: an in vivo MR three-dimensional surface rendering study, 7
Kimura, K., see Tsuzuki, K., 151
Kinsberg, E.R., see Clark, R.B., 255
Kirkbride, V., see Cady, E.B., 201
Kishi, E., see Takahashi, A., 235
Kiuchi, K., see Nagatsu, T., 44
Kobayashi, K., see Nagatsu, T., 44
Kondo, H., see Tanaka, O., 17
Konishi, Y., Kamegai, M., Takahashi, K., Kunishita, T. and Tabira, T., Production of interleukin-3 by murine central nervous system neurons, 271
Kosugi, T., see Takechi, H., 279
Koulchitsky, S.V., Azev, O.A., Gourine, A.V. and Kulchitsky, V.A., Capsaicin-sensitive area in the ventral surface of the rat medulla, 129
Kovács, I., see Kardos, J., 73
Krause, J.E., see Ciofi, P., 193
Kristensen, P., see Fink-Jensen, A., 115
Kristensson, K., see Peng, Z.-C., 104
Kristensson, K., see Xu, X.-J., 263
Krivanek, J., see Bielavska, E., 239
Kulchitsky, V.A., see Koulchitsky, S.V., 129
Kunishita, T., see Konishi, Y., 271

Lambert, G.A., see Angus-Leppan, H., 275
Lantos, P.L., see Arai, N., 197
Lavergne, A., Frain, O., Guibert, B., Faucon Biguet, N. and Leviel, V., Regulation of tyrosine hydroxylase gene expression in mesencephalic dopamine neurons: effect of imipramine treatment, 167
Leigh, P.N., see Willis, C.L., 159
Leviel, V., see Lavergne, A., 167
Liao, K., see Mei, L., 21
Lin, L.-F.H., see Hoffer, B.J., 107
Lindström, L., see Mathé, A.A., 138
Liposits, Z., see Hrabovszky, E., 143
Lorek, A., see Cady, E.B., 201
Lovat, P.E., Lowis, S.P., Pearson, A.D.J., Malcolm, A.J. and Redfern, C.P.F., Concentration-dependent effects of 9-*cis* retinoic acid on neuroblastoma differentiation and proliferation in vitro, 29
Lowis, S.P., see Lovat, P.E., 29

Macaya, A., see Ferrer, I., 77
Macq, A.-F., Philippe, B. and Octave, J.-N., The amyloid peptide of Alzheimer's disease is not produced by internal initiation of translation generating C-terminal amyloidogenic fragments of its precursor, 227
Malcolm, A.J., see Lovat, P.E., 29
Malz, C.R., see Feyerabend, B., 66
Mantle, D., see Ince, P.G., 87
Markesbery, W.R., see Ince, P.G., 87
Marriott, D.R., see Too, H.-P., 185
Martin, D., see Hoffer, B.J., 107
Martín, F., see Ferrer, I., 77
Maruyama, Y., see Takahashi, A., 235
Mathé, A.A., Ågren, H., Lindström, L. and Theodorsson, E., Increased concentration of calcitonin gene-related peptide in cerebrospinal fluid of depressed patients. A possible trait marker of major depressive disorder, 138
Mathiau, P., Bakalara, N. and Aubineau, P., Tryptophan hydroxylase can be present in mast cells and nerve fibers of the rat dura mater but only mast cells contain serotonin, 133
Matsuo, K., see Ogawa, S., 123
Mazzuca, M., see Ciofi, P., 193
McCarley, R.W., see Kikinis, R., 7
Mei, L., Liao, K. and Huganir, R.L., Characterization of substrate specificity of the protein tyrosine phosphatase purified from the electric organ of *Torpedo californica*, 21
Meldrum, B.S., see Willis, C.L., 159
Menegazzi, M., Carcereri De Prati, A. and Grassi Zucconi, G., Differential expression pattern of *jun B* and *c-jun* in the rat brain during the 24-h cycle, 295
Meyer, D.L., see Feyerabend, B., 66
Miki, M., see Hisano, S., 80
Misgeld, U., see Rohrbacher, J., 95
Mishina, M., see Hirano, T., 172
Mizuguchi, T., see Nagatsu, T., 44
Mori, S., see Iwakiri, H., 83
Mori, S., see Takechi, H., 279
Morita, S., see Nagatsu, T., 44
Mormède, P., see Chaouloff, F., 217
Moroji, T., see Baba, A., 119
Mu, X.S., see Yang, K., 287
Mu, X.S., see Yang, K., 291

Nagao, T., see Barbarosie, M.T., 208
Nagatsu, T., Nakahara, D., Kobayashi, K., Morita, S., Sawada, H., Mizuguchi, T. and Kiuchi, K., Peripherally administered (6*R*)-tetrahydrobiopterin increases in vivo tyrosine hydroxylase activity in the striatum measured by microdialysis both in normal mice and in transgenic mice carrying human tyrosine hydroxylase, 44
Nakahara, D., see Nagatsu, T., 44
Naukkarinen, H., see Grönroos, M., 163
Nguyen, C., see Zochodne, D.W., 3
Nishiyama, S., see Takechi, H., 279
Norenberg, M.D., see Itzhak, Y., 251
Nunn, P.B., see Willis, C.L., 159

O'Donnell, B.F., see Kikinis, R., 7
Octave, J.-N., see Macq, A.-F., 227
Ogawa, S., Katsuragi, T., Matsuo, K. and Furukawa, T., Inhibitory effect of okadaic acid on noradrenaline exocytosis from guinea-pig vas deferens (Neuroscience Letters 178 (1994) 144–146 (NSL 10838))., 123
Ohshima, K., see Hiramatsu, K., 37
Oka, T., see Iwakiri, H., 83
Okada, H., see Takechi, H., 279
Okada, Y., see Fujiwara, F., 283
Okada, Y., see Hirai, H., 33
Okamura, H., Tanaka, M., Kanemasa, K., Ban, Y., Inouye, S.-I.T. and Ibata, Y., In situ hybridization histochemistry of *vghm1f* mRNA in the rat suprachiasmatic nucleus: co-localization with vasopressin/neurophysin and VIP/PHI, 181
Olausson, B., see Angus-Leppan, H., 275
Olsson, T., see Xu, X.-J., 263
Onoe, H., see Takechi, H., 279
Onoe, K., see Takechi, H., 279
Osmund, E., see Cady, E.B., 201
Osuna, C., see Harmouch, A., 303

Otten, U., see Gadient, R.A., 243
Owen-Reece, H., see Cady, E.B., 201

Papp, M.I., see Arai, N., 197
Parent, A., see Fortin, M., 51
Pastor, C., see Clos, M.V., 125
Pearson, A.D.J., see Lovat, P.E., 29
Peng, Z.-C., Kristensson, K. and Bentivoglio, M., Dysregulation of photic induction of Fos-related protein in the biological clock during experimental trypanosomiasis, 104
Penrice, J., see Cady, E.B., 201
Perez-Polo, J.R., see Taglialatela, G., 231
Perez-Polo, J.R., see Yang, K., 291
Perez-Polo, J.R., see Yang, K., 287
Pericak-Vance, M.A., see Huang, D.Y., 55
Perry, R., see Hardy, J., 1
Pertovaara, A., see Grönroos, M., 163
Philippe, B., see Macq, A.-F., 227
Pietrzyk, U., see Herholz, K., 47
Planas, A., see Ferrer, I., 77
Planas, A.M., see Soriano, M.A., 205
Pórszász, R. and Szolcsányi, J., Antidromic vasodilatation in the striated muscle and its sensitivity to resiniferatoxin in the rat, 267
Price, C., see Démonet, J.-F., 25
Prihar, G., see Hardy, J., 1
Prins, G.S., see Ciofi, P., 193
Prior, P., see Gass, P., 188

Raghavan, R., see Hardy, J., 1
Rakić, L., see Ćulić, M., 259
Rao, V.L.R. and Butterworth, R.F., Alterations of [^{3}H]8-OH-DPAT and [^{3}H]ketanserin binding sites in autopsied brain tissue from cirrhotic patients with hepatic encephalopathy, 69
Redfern, C.P.F., see Lovat, P.E., 29
Reynolds, R., see Cady, E.B., 201
Roberts, G., see Hardy, J., 1
Rodríguez-Farré, E., see Soriano, M.A., 205
Rohrbacher, J., Bijak, M. and Misgeld, U., Suppression by memantine and amantadine of synaptic excitation intrastriatally evoked in rat neostriatal slices, 95
Rosenthal, A., see Davies, A.M., 112
Roses, A.D., see Huang, D.Y., 55

Sabriá, J., see Clos, M.V., 125
Saitoh, T., see Hansen, L.A., 63
Samuel, W., see Hansen, L.A., 63
Šaponjić, J., see Ćulić, M., 259
Saunders, A.M., see Huang, D.Y., 55
Sawada, H., see Nagatsu, T., 44
Schmechel, D.E., see Huang, D.Y., 55
Schröder, H., see Gass, P., 188
Seipp, S. and Buselmaier, W., Isolation of glyceraldehyde 3-phosphate dehydrogenase (Gapdh) cDNA from the distal half of mouse chromosome 16: further indication of a link between Alzheimer's disease and glycolysis, 91
Sermet, E., Gay, N., Baubet, V., Giaume, M., Dalery, J. and Bobillier, P., Triiodothyronine does not affect the average incorporation of L-[^{35}S]methionine in rat brain structures, 213
Serrano, T., see Ferrer, I., 77
Sharkey, K.A., see Zochodne, D.W., 3
Shaw, P.J., see Ince, P.G., 87
Shenton, M.E., see Kikinis, R., 7
Shibata, S., Tominaga, K. and Watanabe, S., Effects of naloxone, morphine and κ-opioid receptor agonists on hypoxia/hypoglycemia-induced reduction of 2-deoxyglucose uptake in hippocampal slices from U-50,488H-tolerant rats, 155
Shima, K., see Tsuzuki, K., 151
Shinozuka, K., see Hirano, T., 172
Soriano, M.A., Planas, A.M., Rodríguez-Farré, E. and Ferrer, I., Early 72-kDa heat shock protein induction in microglial cells following focal ischemia in the rat brain, 205
Strada, O., see Villares, J., 59
Strittmatter, W.J., see Huang, D.Y., 55
Sun, L.R. and Suszkiw, J.B., Pb^{2+} activates potassium currents in bovine adrenal chromaffin cells, 41
Suszkiw, J.B., see Sun, L.R., 41
Suzuki, T., see Baba, A., 119
Szolcsányi, J., see Pórszász, R., 267

Tabira, T., see Konishi, Y., 271
Taglialatela, G. and Perez-Polo, J.R., Developmental profile of the hypothalamo-pituitary-adrenal axis response to nerve growth factor, 231
Taglialatela, G., see Yang, K., 287
Taglialatela, G., see Yang, K., 291
Takahashi, A., Ishimaru, H., Ikarashi, Y., Kishi, E. and Maruyama, Y., Increase of hypothalamic cholinergic activity in 2-deoxyglucose hyperglycemia, 235
Takahashi, K., see Konishi, Y., 271
Takahata, N., see Tsuzuki, K., 151
Takamaru, Y., see Tsuzuki, K., 151
Takechi, H., Onoe, H., Imamura, K., Onoe, K., Kakiuchi, T., Nishiyama, S., Yoshikawa, E., Mori, S., Kosugi, T., Okada, H., Tsukada, H. and Watanabe, Y., Brain activation study by use of positron emission tomography in unanesthetized monkeys, 279
Tamaki, N., see Fujiwara, F., 283
Tanaka, M., see Okamura, H., 181
Tanaka, O. and Kondo, H., Localization of mRNAs for three novel members (β3, β4 and γ2) of phospholipase C family in mature rat brain, 17
Tandon, L., see Ince, P.G., 87
Theodorsson, E., see Mathé, A.A., 138
Tominaga, K., see Shibata, S., 155
Too, H.-P., Marriott, D.R. and Wilkin, G.P., Preprotachykinin-A and substance P receptor (NK1) gene expression in rat astrocytes in vitro, 185
Tortosa, A., see Ferrer, I., 77
Tsukada, H., see Takechi, H., 279
Tsuzuki, K., Fukatsu, R., Takamaru, Y., Kimura, K., Abe, M., Shima, K., Fujii, N. and Takahata, N., Immunohistochemical evidence for amyloid β in rat soleus muscle in chloroquine-induced myopathy, 151

Uemura, N., see Hisano, S., 80
Urban, N.N., see Chinestra, P., 177

van der Meide, P.H., see Xu, X.-J., 263
Villares, J., Strada, O., Faucheux, B., Javoy-Agid, F., Agid, Y. and Hirsch, E.C., Loss of striatal high affinity NGF binding sites in progressive supranuclear palsy but not in Parkinson's disease, 59

Watanabe, S., see Shibata, S., 155
Watanabe, Y., see Takechi, H., 279
Weisgraber, K.H., see Huang, D.Y., 55